

HEPATITIS C VIRUS SPECIALTY CARE PROGRAM

Phone: **877-324-2501** • Fax: **888-972-4110**



©2017 KloudScript, Inc. - All rights reserved.

1 PATIENT INFORMATION: Name:			2 PRESCRIBER INFORMATION: Name:			
Address:						
City:			State: Z			
Phone:	•	•	Fax:	•		
Email:			DEA:			
	_		Phone:			
STATEMENT OF MEI						
Diagnostic Information			Labs			
Date of Diagnosis: ICD-		ce:	ALT: HGB:			
Genotype: Subtype: _	Q80K: 🗖 Positive	☐ Negative (For Genotype 1a)	AST: HCV RNA			
Indicate Patient Status: ☐ Naïve ☐ Partial Responder ☐ Non-responder ☐ Null-responder ☐ Relapser			PLT: SrCr:			
Duration of Previous Therapy: Weeks From: To: To: Cirrhosis: ☐ No ☐ Yes If Yes: ☐ Compensated ☐ Decompensated			NS5A Resistance Assay: Date:			
History of Liver Biopsy? ☐ No ☐ Yes If Yes, Please Attach Results			NSSA Resistance Assay.	Jaie:		
□ Fibrosure or □ Fibroscan: Results:			Medication List and Contra	indication	าร	
Extra-Hepatic Manifestations: ☐ Ascites ☐ Hepatic Encephalopathy ☐ Thrombocytopenia ☐ Other: Does the patient need liver transplantation? ☐ Yes ☐ No			☐ Attach Medication List			
HBsAg and anti-HBc Test: Positive Negative Pate:			Is the patient interferon ineligible? ☐ No ☐ Yes			
If Prior Authorization is denied, recommended formulary alternatives will be provided to the prescriber based upon the patient's insurance coverage.			☐ Anxiety ☐ Depression ☐ Pulmonary Abnormalities			
			Renal Insufficiency Other:			
_			g O Patient Trained in MD Office O Manuf			
5 PICK UP OR DELIVE	RY: O Delivery to Patient	t's Home O Delivery to I	Physician's Office O Pharmacy t	o Coordin	nate	
6 INSURANCE INFORM						
PRESCRIPTION: Duration of Patient Name:	of Therapy: 🗆 8 weeks 😐 12		er itient's Date of Birth:			
Medication	Dosage & Strength		Direction	QTY I	Refills	
	□ 30mg Tablets	with or without Ribavirin	vithout food in combination with Sovaldi®	28		
□ DAKLINZA™	☐ 60mg Tablets	with or without Ribavirin	vithout food in combination with Sovaldi®	28		
	□ 90mg Tablets	☐ Take 90mg daily with or w with or without Ribavirin	□ Take 90mg daily with or without food in combination with Sovaldi® with or without Ribavirin			
□ EPCLUSA®	☐ 400/100mg Tablets	☐ Take one tablet daily with	or without food	28		
☐ HARVONI®	□ 90/400mg Tablets	☐ Take one tablet daily with	or without food	28		
□ MAVYRET™	□ 100/40mg Tablet	☐ Take three tablets orally o	nce daily with food	1 Carton		
□ OLYSIO®	☐ 150mg Capsules	☐ Take one 150mg capsule	orally once a day	28		
☐ SOVALDI®	☐ 400mg Tablets	☐ Take one 400mg tablet or	☐ Take one 400mg tablet orally once a day			
□ VOSEVI®	☐ 400/100/100mg Tablets	☐ Take one tablet orally onc	e daily with food	28		
☐ MODERIBA Dose Pack™	☐ 600mg per day		morning/400mg tablet every evening			
☐ RIBASPHERE Riba Pack®	■ 800mg per day■ 1000mg per day		morning/400mg tablet every evening morning/400mg tablet every evening			
The for the field fact.	☐ 1200mg per day		morning/600mg tablet every evening			
□ MODERIBA [™]	□ 200mg Tablets	□ Take tablets/	capsules every morning and,			
☐ RIBASPHERE®☐ RIBAVIRIN	□ 200mg Capsules	☐ Take tablets/	capsules every evening			
	□ EE0ma Tablata	□ Taka ana tablat twica dail	ith as without food	60		
☐ XIFAXAN® ☐ ZEPATIER®	☐ 550mg Tablets	☐ Take one tablet daily with	•	60		
	□ 50/100mg Tablets	☐ Take one tablet daily with	or without lood	28		
PRESCRIBER SIGNAT	LIDE.					
					-	
Signature:	Date:			ate:	-	